



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2015

Plant germline formation: common concepts and developmental flexibility in sexual and asexual reproduction

Schmidt, Anja ; Schmid, Marc W ; Grossniklaus, Ueli

Abstract: The life cycle of flowering plants alternates between two heteromorphic generations: a diploid sporophytic generation and a haploid gametophytic generation. During the development of the plant reproductive lineages - the germlines - typically, single sporophytic (somatic) cells in the flower become committed to undergo meiosis. The resulting spores subsequently develop into highly polarized and differentiated haploid gametophytes that harbour the gametes. Recent studies have provided insights into the genetic basis and regulatory programs underlying cell specification and the acquisition of reproductive fate during both sexual reproduction and asexual (apomictic) reproduction. As we review here, these recent advances emphasize the importance of transcriptional, translational and post-transcriptional regulation, and the role of epigenetic regulatory pathways and hormonal activity.

DOI: <https://doi.org/10.1242/dev.102103>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-107005>

Journal Article

Published Version

Originally published at:

Schmidt, Anja; Schmid, Marc W; Grossniklaus, Ueli (2015). Plant germline formation: common concepts and developmental flexibility in sexual and asexual reproduction. *Development*, 142(2):229-241.

DOI: <https://doi.org/10.1242/dev.102103>

REVIEW

Plant germline formation: common concepts and developmental flexibility in sexual and asexual reproduction

Anja Schmidt*, Marc W. Schmid and Ueli Grossniklaus*

ABSTRACT

The life cycle of flowering plants alternates between two heteromorphic generations: a diploid sporophytic generation and a haploid gametophytic generation. During the development of the plant reproductive lineages – the germlines – typically, single sporophytic (somatic) cells in the flower become committed to undergo meiosis. The resulting spores subsequently develop into highly polarized and differentiated haploid gametophytes that harbour the gametes. Recent studies have provided insights into the genetic basis and regulatory programs underlying cell specification and the acquisition of reproductive fate during both sexual reproduction and asexual (apomictic) reproduction. As we review here, these recent advances emphasize the importance of transcriptional, translational and post-transcriptional regulation, and the role of epigenetic regulatory pathways and hormonal activity.

KEY WORDS: Cell fate acquisition, Gene regulation, Germline development, Plant reproduction, Polarity

Introduction

In higher plants, diverse and versatile strategies have evolved to ensure reproductive success. During gametogenesis (see Glossary, Box 1), the male (pollen) and female (embryo sac) gametophytes, which harbour the male (sperm) and female (egg and central cell; see Glossary, Box 1) gametes, respectively, form in specialized reproductive tissues of the flower: the anther and ovule (Fig. 1). The multicellular gametophytes are formed following meiosis of spore mother cells (see Glossary, Box 1), thus producing reduced gametes that harbour half the chromosome number of the maternal sporophyte (haploid in case of diploid plants). During sexual reproduction (see Glossary, Box 1), sperm cells fuse with both the egg and the central cell in the process of double fertilization, giving rise to the embryo and endosperm, respectively, the major components of the seed (Fig. 1). The embryo constitutes the next sporophytic generation, while the endosperm is a terminal nourishing tissue for the embryo and also provides the majority of calories for human and animal consumption. Haploid plants can also form directly from male and female gametes. While this process occurs at very low frequencies in nature, it can be induced in culture and by mutation, and hence is being exploited to accelerate plant breeding (Germanà, 2011). By contrast, during vegetative reproduction (see Glossary, Box 1) and somatic embryogenesis (see Glossary, Box 1), which are two distinct types of asexual reproduction (see Glossary, Box 1), new plants develop without the formation of gametes and seeds. However, plants can also produce seeds via asexual reproduction, avoiding the need for

fertilization, in a process known as gametophytic apomixis (hereafter referred to as apomixis, see Glossary, Box 1). Apomixis occurs in more than 400 plant species belonging to ~40 genera.

Both sexual reproduction and apomixis have distinct advantages for natural plant populations and agricultural applications. Sexual reproduction leads to genetically and phenotypically variable offspring, thus forming the basis for plant adaptation to changing environments and allowing for the breeding of new varieties. By contrast, apomixis produces clonal offspring that are genetically identical to the mother plant, thus fixing complex genotypes. Although apomixis is rare among crop plants, the engineering of apomictic crops promises great potential and economical value for crop production and for other applications in agriculture (Koltunow et al., 1995; Vielle-Calzada et al., 1996; Grossniklaus et al., 1998a,b; Spillane et al., 2004).

Over the past decade, plant sexual and apomictic germline formation has attracted the attention of scientists for a number of reasons: (1) the transition from sporophytic to reproductive fate by

Box 1. Glossary

Apomeiosis. The omission or abortion of meiosis during sporogenesis

Apomictic initial cell (AIC). The first cell in the apomictic female germline that omits or aborts meiosis

Apomixis. Asexual reproduction via seed formation

Apospory. The formation of an unreduced female gametophyte from an apomictic initial cell (AIC) developing adjacent to the sexual germline in the ovule

Archisporial cell. The cell giving rise (with or without division) to the spore mother cell

Asexual reproduction. Reproduction without the fusion of gametes

Central cell. The female gamete giving rise to the endosperm

Diplospory. The apomeiotic formation of an unreduced female gametophyte from an AIC at the position of the megaspore mother cell

Egg cell. The female gamete giving rise to the embryo

Functional megaspore (FMS). The cell that develops into the female gametophyte

Gametogenesis. The development of gametophytes from spores

Parthenogenesis. The formation of an embryo from an unfertilized egg cell

Pseudogamy. The fertilization-dependent formation of endosperm from a central cell in apomicts

Sexual reproduction. The mode of reproduction whereby female (egg) and male (sperm) gametes fuse to form a zygote

Somatic embryogenesis. The formation of an embryo from a sporophytic cell without gamete and seed formation

Sporogenesis. The formation of spores from spore mother cells

Spore mother cell. The first cell of the reproductive lineage, formed from sporophytic cells in female and male reproductive tissues of the flower

Synergid cells. Accessory cells of the mature female gametophyte that are important for pollen tube guidance and reception

Vegetative reproduction. A form of reproduction in which a new plant is formed without the formation of an embryo

Institute of Plant Biology and Zürich-Basel Plant Science Centre, University of Zürich, Zollikerstrasse 107, Zürich CH-8008, Switzerland.

*Authors for correspondence (aschmidt@botinst.uzh.ch; grossnik@botinst.uzh.ch)




Fig 1. The life cycle of a plant. Plants have a more complex life cycle than animals, alternating between two heteromorphic generations: the sporophyte and the gametophyte. In the diploid sporophyte, distinct cells undergo meiosis and produce haploid spores. These give rise to multicellular haploid gametophytes, which produce gametes through mitotic divisions. The fusion of a male (sperm) and a female (egg) gamete results in the formation of a zygote, which constitutes the sporophyte. The example depicted, *Hieracium pilosella*, follows the common life cycle of angiosperms. The anthers of the flower produce the male gametophyte (called the pollen or microgametophyte), which consists of three cells: two sperm cells (SC) and one vegetative cell (VC). The female gametophyte (called the embryo sac or megagametophyte) is embedded in maternal sporophytic tissues of the ovule. The latter is enclosed in the carpels of the flower. For sexual reproduction, the pollen needs to germinate on the stigma and to deliver the two sperm cells to the female gametophyte. Fertilization, the transition from the gametophytic to the sporophytic generation, occurs within the ovule. In addition to sexual reproduction, plants can frequently reproduce asexually, e.g. by stolon outgrowth (vegetative reproduction) or via the formation of calli in culture, followed by somatic embryogenesis and development into an adult plant. CC, central cell; EGG, egg cell; SC, sperm cell; SYN, synergid cell; VC, vegetative cell.

manipulation of plant reproduction for agricultural use and crop improvement. Accordingly, many studies have focussed on determining the gene expression profiles, epigenetic mechanisms and regulatory pathways involved in germline development (reviewed by Drews and Koltunow, 2011; Sprunck and Gross-Hardt, 2011; Schmidt et al., 2012; Gutierrez-Marcos and Dickinson, 2012; Wüest et al., 2013). Here, we focus on recent studies that have elucidated the molecular mechanisms underlying the acquisition of reproductive fate in sexual and apomictic species, the determination of meiosis versus apomeiosis (see Glossary, Box 1), and the polar development of the female gametophyte.

Development of the plant reproductive lineages

The formation of the plant reproductive lineages proceeds in two distinct phases: during sporogenesis (Glossary, Box 1), spores are formed by sporophytic (somatic) cells, whereas during gametogenesis the spores develop into mature gametophytes that harbour the male or female gametes (Fig. 2). During the course of evolution, the gametophytic phase of the plant life cycle, which is dominant in bryophytes (i.e. liverworts, hornworts and mosses), has been dramatically reduced to only a few cells in the angiosperms (flowering plants). Thus, unlike in most animals, where the germline is set aside early in embryogenesis, the plant germline is determined only late in development, during floral organ formation. Here, we consider the spore mother cells to be the first cells of the germline, as the lineage of the gametes can unambiguously be traced back to them (Grossniklaus, 2011). However, it should be noted that, because gametophytes consist of both gametic and non-gametic accessory cells and the germline is defined as the cell lineage that differentiates into gametes, some authors place the determination of the germline later during gametophyte development to the immediate precursors of the gametes (e.g. Berger and Twell, 2011; Twell, 2011).

The formation of the male reproductive lineage begins with the differentiation of a microspore mother cell (MiMC) in the developing anthers; the periclinal division of archesporial cells (see Glossary, Box 1) gives rise to outer parietal cells and inner sporogenous cells, and the MiMCs differentiate from the latter. The MiMC undergoes meiosis to give rise to a tetrad of microspores (Fig. 2), each of which undergoes an asymmetric division (termed pollen mitosis I, PMI) to form a vegetative and a generative cell (Borg et al., 2009). During pollen mitosis II (PMII), the generative cell forms two sperm cells (male gametes), while the vegetative cell does not divide again. The

reprogramming a somatic cell is a key step in the plant life cycle; (2) during gametogenesis, a few rounds of mitosis and cellularization lead to the formation of functionally distinct cell types that are all derived from a single spore, a process ideally suited to address fundamental questions in developmental biology; and (3) understanding the molecular mechanisms that determine sexual or asexual fate decisions is a precondition for the targeted

sperm cells are then delivered to the female gametes by the pollen tube, which forms via growth of the vegetative cell. The timing of PMII varies in different species; in most plant species, PMII takes place in the growing pollen tube but in some species, including *Arabidopsis* and maize, the generative cell divides before the pollen is released from the anther (Boavida et al., 2005).

During formation of the female sexual reproductive lineage, typically a single somatic cell per ovule acquires reproductive fate and differentiates to form an archesporial cell. It can be distinguished from the surrounding cells by its subepidermal localization and its enlarged size. In the sexual model species *Arabidopsis*, as in most species, the archesporial cell directly differentiates into a megaspore mother cell (MMC) without intervening divisions. The MMC is defined by its commitment to the meiotic fate and gives rise to a tetrad of megaspores (Fig. 2). Typically, only one functional megaspore (FMS; Glossary, Box 1) survives while the others degenerate (Fig. 2). Interestingly, the FMS occupies a defined position in the ovule, suggesting that this is important for its survival and cell fate acquisition. A role for signalling from sporophytic ovule tissues during the selection of the FMS has been discussed (Koltunow, 1993; Grossniklaus and Schneitz, 1998; Koltunow and Grossniklaus, 2003) and, in maize, the accumulation of callose in the cell walls of the degenerating megaspores has been hypothesized to play a role in shielding these cells from such signals (Russell, 1979). The FMS, in turn, typically undergoes three mitotic divisions to form a syncytial female gametophyte (Fig. 2). In most species, cellularization results in an eight-nucleate, seven-celled mature gametophyte (embryo sac), referred to as a *Polygonum* type embryo sac. It harbours the two female gametes, the synergid cells (see Glossary, Box 1), which are important for pollen tube guidance and reception, and three antipodal cells (Fig. 2). Although the role of the antipodal cells remains unclear, they might be involved in transferring nutrients from the surrounding sporophytic tissues to the embryo sac (Raghavan, 1997). The *Polygonum* type embryo sac occurs in ~70% of all angiosperms, including the model systems *Arabidopsis thaliana* (mouse ear cress), *Zea mays* (maize) and *Oryza sativa* (rice), and many apomictic species (Drews and Koltunow, 2011).

From the beginning of its development, the female gametophyte is highly polarized, suggesting that positional information may play a role in cell fate acquisition (Grossniklaus and Schneitz, 1998; Lituiev and Grossniklaus, 2014). The exact position of nuclei within the syncytium may thus be an important factor for cell specification during cellularization (Sundaresan and Alandete-Saez, 2010; Sprunck and Gross-Hardt, 2011). It is also evident that variations in this developmental pattern exist: while megasporogenesis typically leads to a single surviving one-nucleate FMS (monosporic megasporogenesis), failures in cell plate formation after meiosis I or after both meiotic divisions can lead to two- or four-nucleate FMSs, developmental patterns referred to as bisporic or tetrasporic megasporogenesis, respectively (Maheshwari, 1950; Willemse and Went, 1984; Haig, 1990; Huang and Russell, 1992; Drews and Koltunow, 2011). Other developmental variations concern the number of mitoses during megagametogenesis before cellularization, the possibility of additional mitoses after cellularization, and the timing of the fusion of the polar nuclei in the central cell (Maheshwari, 1950; Drews and Koltunow, 2011).

Sexual reproduction and apomixis are interrelated

Compared with sexual reproduction, apomixis differs only in three key developmental steps (Fig. 2). First, female meiosis is circumvented, in a process referred to as apomeiosis, leading to

the formation of unreduced megaspores and, consequently, unreduced female gametes. The first cell of the apomictic lineage is termed an apomictic initial cell (AIC; see Glossary, Box 1). The AIC is either formed at the position of the MMC and omits or aborts meiosis (diplospory; see Glossary, Box 1) to give rise to an unreduced FMS, or is derived from a somatic cell in close proximity to the MMC that directly differentiates into an unreduced FMS (apospory; see Glossary, Box 1) (Bicknell and Koltunow, 2004). Usually, male meiosis is unaffected but unreduced pollen can also be produced in some apomicts (Bicknell and Koltunow, 2004). Second, the egg develops into an embryo in the absence of fertilization in a process known as parthenogenesis (see Glossary, Box 1). Currently, the molecular mechanisms that activate the egg cell and initiate embryogenesis are unknown. Third, the central cell can form endosperm either autonomously or after fertilization (pseudogamy; see Glossary, Box 1). Functional endosperm formation in pseudogamous apomicts requires adaptations in either megagametogenesis (the production of a four-nucleate embryo sac), microgametogenesis (the formation of unreduced sperm cells) or double fertilization to ensure a balanced endosperm with the correct 2:1 ratio of maternal to paternal genomes crucial for seed development in many species (Grossniklaus, 2001; Koltunow and Grossniklaus, 2003; Spillane et al., 2004). For example, in maize *indeterminant gametophyte1* (*igl1*) mutants, abnormal numbers of nuclei are formed in the female gametophyte, leading to an aberrant maternal to paternal genome ratio in the endosperm, which results in seed abortion (Lin, 1984; Huang and Sheridan, 1996). In autonomous apomicts, the requirement for a balanced endosperm is alleviated, likely also depending on specific adaptations that are under genetic control.

The acquisition and restriction of reproductive fate

During sexual reproduction, only one somatic cell per ovule is usually committed to the reproductive fate. However, it is not fully understood what determines the commitment of this somatic cell to initiate germline development and what prevents the formation of additional germline cells in the same ovule. In higher plants, it has been hypothesized that the MMC represses the formation of additional MMCs and thus restricts germline formation to only one cell per ovule (Grossniklaus and Schneitz, 1998). In support of the hypothesis that the germline itself suppresses the formation of additional germline lineages, the formation of multiple female gametophytes per ovule has been described in *Trimenia*, an ancient angiosperm taxon, where tip growing female gametophytes compete to reach the site of fertilization (Bachelier and Friedman, 2011).

Initial insights into the signalling pathways that regulate the restriction of germline fate came from the analyses of mutants in maize, rice and *Arabidopsis* (summarised in Table 1). In rice carrying mutations in *MULTIPLE SPOROCTE* (*MSP1*) and in *Arabidopsis* plants carrying mutations in the orthologue *EXTRA SPOROGENOUS CELLS/EXCESS MICROSPOROCTES1* (*EXS/EMS1*) or mutations in *SOMATIC EMBRYOGENESIS RECEPTOR KINASE1* and 2 (*SERK1/2*), more MiMCs develop per anther in comparison to the wild type (Canales et al., 2002; Zhao et al., 2002; Nonomura et al., 2003; Albrecht et al., 2005; Colcombet et al., 2005; Jia et al., 2008). These genes encode leucine-rich receptor kinases (*MSP1* and *EXS/EMS1*) and LRR receptor-like serine threonine kinases (*SERK1/2*) (Canales et al., 2002; Zhao et al., 2002; Nonomura et al., 2003; Albrecht et al., 2005; Colcombet et al., 2005; Jia et al., 2008). Similar phenotypes have been described in mutant in which the genes *TAPETUM DETERMINANT1* (*TPD1*) in




Fig. 2. Male and female gametophyte development in sexually reproducing and apomictic plant species. Germline development starts with the differentiation of sporophytic cells into spore mother cells (female, megaspore mother cell, MMC; male, microspore mother cell, MiMC) that, in sexually reproducing species undergo meiosis to give rise to four haploid spores. During male gametogenesis, the four spores separate and form unicellular microspores (MS), which grow by cell expansion to form unicellular pollen (1cP). The first asymmetric mitosis produces bicellular pollen (e2cP) containing a large vegetative cell (VC) and a small generative cell (GC). The GC detaches from the cell wall and becomes engulfed by the VC. Sperm cells (SC) are formed during the second mitosis of the GC. The mature pollen (mP) consists of a VC, which will form the pollen tube, and two SCs that mediate double fertilization. During female gametogenesis, three of the four spores degenerate, leaving one functional megaspore (FMS), which undergoes three mitotic divisions in a syncytium to give rise to the early/late two-nucleate, four-nucleate and then eight-nucleate embryo sac (e/2nES, 4nES, 8nES). Nuclear migration and concomitant cellularization eventually lead to the formation of a mature embryo sac (mES), a highly polarized structure that contains four distinct cell types: two synergid cells (SYN), the egg cell (EGG), the central cell (CC) and antipodal cells (APs), which degenerate prior to fertilization. In apomictic species, different mechanisms can lead to the formation of unreduced gametes. In diplospory, the apomictic initial cell (AIC) may initiate meiosis but restitution results in the formation of two unreduced AICs, one of which degenerates. By contrast, aposporous apomicts form a FMS-like cell at a different position in the ovule. The unreduced AICs then develop into unreduced female gametophytes. Meiosis on the male side is usually normal in apomicts. Female gametophyte stages (FG) are according to Christensen et al. (1997). p, precursor of.

Arabidopsis, *MULTIPLE ARCHESPORIAL CELLS (MAC1)* in maize and its rice orthologue *OsTDL1A* are disrupted. These genes encode small secreted proteins identified as the putative ligands of the MSP1 or EXS/EMS1 receptor kinases (Sheridan et al., 1996, 1999; Yang et al., 2003, 2005; Zhao et al., 2008; Wang et al., 2012; Kelliher and Walbot, 2012). Unlike in *Arabidopsis*, in rice and maize this pathway also affects female sporogenesis, indicating differences in the mechanism of repression of additional sporocytes (Sheridan et al., 1996; Zhao et al., 2008; Nonomura et al., 2003). In maize and rice, excess archesporial cells were observed, leading to the formation of more sporogenous cells (Zhang and Yang, 2014). In *Arabidopsis*, however, the pathway plays a role in cell fate decisions and cell specification after the periclinal division of the archesporial cell. As recently demonstrated, EXS/EMS1 forms complexes with SERK1/2 to control the proliferation of tapetal cells in the anther (Albrecht et al., 2005; Colcombet et al., 2005; Feng and Dickinson, 2010). Interestingly, partially complementary expression patterns have been reported for *TPD1* and *EXS/EMS1*, which are predominantly expressed in sporogenous cells and tapetal cells, respectively, at the developmental stages at which the mutant phenotypes are first established, indicating signalling between cell types (Yang et al., 2003).

In aposporous apomicts, such repression of additional germline lineages is not active, as both an enlarged somatic AIC and the MMC can initiate reproductive lineages (Fig. 1). Formation of the AIC in *Hieracium pilosella* even depends on differentiation and meiosis of the sexual MMC (Koltunow et al., 2011). Thus, as suggested by the signalling pathways described above, communication between cell types during sporogenesis seems to be involved in cell type specification and the acquisition or restriction of germline fate. It remains unclear whether this is achieved by overcoming the mechanism that usually represses the development of additional germline cells or by an alternative signalling pathway that induces reproductive fate in an additional somatic cell, or whether a combination of both of these mechanisms is involved. However, it should be noted that, once established, the apomictic lineage often suppresses the further development of the sexual female gametophyte (Koltunow et al., 2011), suggesting that distinct control mechanisms exist at these developmental steps.

In diplosporous species, the AIC omits or aborts meiosis producing an unreduced FMS (Fig. 2; Bicknell and Koltunow, 2004). As in sexual species, diplosporous apomicts typically develop only one germline lineage per ovule, suggesting that the processes by which gametophytic fate is acquired in apospory and diplospory follow distinct developmental principles. It is unknown whether apomictic fate is regulated by related or alternative molecular mechanisms in diplosporous and aposporous species. Investigations into this question have proved to be technically challenging as the female germline is deeply embedded in maternal floral tissues. Nevertheless, recent methodological advances have allowed the transcriptional profiling of such rare cell types by combining laser-assisted microdissection or micromanipulation with microarray and/or RNA-Seq analyses. These studies have provided novel insights into the transcriptional basis of germline specification and development (Wüest et al., 2010; Schmidt et al., 2011, 2012, 2014; Schmid et al., 2012; Okada et al., 2013; Wüest et al., 2013; Abiko et al., 2013; Anderson et al., 2013; Chettoor et al., 2014).

New insights into the issue of how cell specification is regulated during diplospory, when compared with sexual or aposporous reproduction, were recently provided by cell type-specific transcriptome analyses of the reproductive lineage in *Boechera gunnisoniana*, a diplosporous apomict that is related to sexual *A. thaliana* (Schmidt et al., 2014). Comparative transcriptome analyses detected a number of commonalities between the sexual MMC and the diplosporous AIC (Schmidt et al., 2014). Importantly, significant differences in the activities of a number of regulatory pathways were also observed, including differences in cell cycle regulation, hormonal pathways, signal transduction and epigenetic regulatory pathways (Schmidt et al., 2014). Through comparisons with a transcriptome dataset of the AIC of *Hieracium praealtum* (Okada et al., 2013), this study suggests interesting differences between the regulatory mechanisms specifying a diplosporous or an aposporous AIC (Schmidt et al., 2014). Importantly, the *H. praealtum* AIC seems to have already adopted a gametophytic fate (Okada et al., 2013). In agreement with this acquisition of a FMS fate without meiotic division, a number of meiotic genes are not expressed in the *H. praealtum* AIC (Okada et al., 2013). By contrast, the majority of 25 core meiotic genes are expressed in the AIC of

Table 1. Genes involved in the restriction of additional sporogenous cells during plant reproductive development

Gene	Species	Restricts number of MiMCs, MMCs or both	Type/function of protein encoded	Reference(s)
<i>MSP1</i>	<i>O. sativa</i>	Restricts the number of sporocytes in anther and ovule	Leucine-rich repeat receptor-like kinase; orthologue of <i>Arabidopsis</i> <i>EXS/EMS1</i>	Nonomura et al., 2003
<i>EXS/EMS1</i>	<i>A. thaliana</i>	Restricts the number of microsporocytes	Leucine-rich repeat receptor-like kinase involved in regulating the proliferation of tapetal cells during anther development	Canales et al., 2002; Zhao et al., 2002; Feng and Dickinson, 2010
<i>SERK1/2</i>	<i>A. thaliana</i>	Restricts the number of microsporocytes	Leucine-rich repeat receptor-like kinases; forms complexes with <i>EXS/EMS1</i> in tapetal cells	Albrecht et al., 2005; Colcombet et al., 2005
<i>TPD1</i>	<i>A. thaliana</i>	Restricts the number of microsporocytes	Small secreted protein; interacts with <i>EXS/EMS1</i>	Yang et al., 2003; Yang et al., 2005
<i>MAC1</i>	<i>Z. mays</i>	Restricts the number of archesporial cells in anther and ovule	Small secreted protein; orthologue of <i>OsTDL1A</i>	Sheridan et al., 1996; Sheridan et al., 1999; Wang et al., 2012
<i>OsTDL1A</i>	<i>O. sativa</i>	Restricts the number of sporocytes in anther and ovule	Small secreted protein; putative ligand of <i>MSP1</i>	Zhao et al., 2008

A. thaliana, *Arabidopsis thaliana*; *EXS/EMS1*, *EXTRA SPOROGENOUS CELLS/EXCESS MICROSPOROXYTES1*; *MAC1*, *MULTIPLE ARCHESPORIAL CELLS1*; MiMCs, microspore mother cells; MMCs, megaspore mother cells; *MSP1*, *MULTIPLE SPOROCYTE1*; *O. sativa*, *Oryza sativa*; *OsTDL1A*, orthologue of *MAC1*; *SERK1/2*, *SOMATIC EMBRYOGENESIS RECEPTOR KINASE1* and 2; *TPD1*, *TAPETUM DETERMINANT1*; *Z. mays*, *Zea mays*.

the diplosporous species *B. gunnisoniana* before first division restitution (Schmidt et al., 2014). This supports the notion that diplospory results from a modification of the meiotic pathway in an MMC-like cell, while the aposporous AIC becomes directly determined to a gametophytic fate without prior activation of the meiotic program.

Mutations in meiotic genes can lead to diplospory-like modifications of meiosis

Over recent years, investigations into the regulatory processes governing meiosis have allowed the identification of meiotic mutants that generate unreduced gametes (Table 2) (Brownfield and Köhler, 2011; Crismani et al., 2013). For example, mutations in the gene encoding *DYAD/SWITCH1 (SWI1)* lead to apomeiosis and to the formation of rare triploid offspring that retain full parental heterozygosity (Ravi et al., 2008). In *MiMe-1* and *MiMe-2* triple mutants, a diplospory-like division also leads to the formation of unreduced gametes. *MiMe-1* and *MiMe-2* are combinations of *sporulation11-1 (spo11-1)*, *omission of second division1 (osd1)* and *recombination8 (rec8)*, and *spo11-1*, *osd1* and *cyc1;2/tardy asynchronous meiosis (tam)*, respectively (d'Erfurth et al., 2009, 2010). Using these *Arabidopsis* mutants, synthetic clonal seeds have been produced by manipulating the expression of the centromere-specific histone 3 variant CENH3, which leads to paternal genome elimination, in the *dyad/swi1* or *MiMe* mutant background (Marimuthu et al., 2011).

Meiosis and the acquisition of germline fate are affected by abiotic and oxidative stress

Although these mutations in meiotic genes lead to a deregulation of the meiotic program and, eventually, to a switch to a diplospory-like process, little is known about the control of diplospory and meiotic restitution in natural apomicts. Interestingly, abiotic stress can lead to alterations in meiotic cell division (de Storme and Geelen, 2014). For example, in rose (*Rosa spp.*) short periods of heat stress result in partial restitution of male meiosis and the formation of unreduced dyads, but also triads and polyads (Pecirix et al., 2011).

Analysis of the effect of redox status on germline specification in maize revealed another link to abiotic stress. In anthers, germ cell

formation is stimulated by a low oxygen environment or by a low abundance of reactive oxygen species (ROS), which accumulate under different kinds of stress (Kelliher and Walbot, 2012, 2014). This led to the conclusion that reduced oxygen concentration promotes the acquisition of meiotic fate in maize (Kelliher and Walbot, 2012). Contrasting the idea that meiotic fate is acquired under low ROS levels, a recent hypothesis postulates that the evolution and maintenance of meiosis depends on stress and elevated ROS levels (Hörandl and Hadacek, 2013, see Box 2).

In conclusion, although abiotic and oxidative stresses seems to play a role in the transition from somatic to reproductive fate and the regulation of (apo)meiosis, their potential role as a driving force promoting sexual or asexual reproduction remains unclear and warrants further investigation.

Epigenetic regulatory pathways are important for germline specification and the control of sexual versus apomictic reproduction

Disturbance of the meiotic programme typically results in sterility or the diplospory-like formation of unreduced gametes. However, phenotypes resembling apospory or diplospory have also been observed in mutants perturbing epigenetic regulatory pathways, in particular those involving DNA methylation and small RNA-based gene regulation (Olmedo-Monfil et al., 2010; Garcia-Aguilar et al., 2010; Singh et al., 2011).

Epigenetic regulation is involved in a variety of developmental and cell fate decisions by controlling gene activity through DNA or chromatin modifications. For example, ARGONAUTE (AGO) proteins are involved in gene regulation mediated by small RNAs such as microRNAs (miRNA), small interfering RNAs (siRNA) and PIWI-associated RNAs (piRNA) (Meister, 2013). In *Arabidopsis*, 10 AGO proteins have been identified, and these can be grouped into three major clades: the AGO1, AGO5 and AGO10 clade; the AGO2, AGO3 and AGO7 clade; and the AGO4, AGO6, AGO8 and AGO9 clade (Mallory and Vaucheret, 2010). These different clades of AGO proteins engage in different small RNA pathways, with the AGO9 clade being active in the siRNA heterochromatin pathway that regulates the transcriptional silencing of transposons and repeats by mediating DNA methylation and heterochromatin formation (Mallory and Vaucheret, 2010).

Table 2. Mutations that lead to the formation of unreduced female gametophytes by an apospory- or diplospory-like mechanism

Mutation	Species	Description	Type of apomeiosis	Reference(s)
<i>dyad/swi1</i>	<i>A. thaliana</i>	Mutation in core meiotic gene	Diplospory like	Ravi et al., 2008
<i>MiMe-1 (spo11-1, osd1 and rec8)</i>	<i>A. thaliana</i>	Triple mutant of core meiotic genes	Diplospory like	d'Erfurth et al., 2009
<i>MiMe-2 (spo11-1, osd1 and tam)</i>	<i>A. thaliana</i>	Triple mutant of core meiotic genes	Diplospory like	d'Erfurth et al., 2010
<i>ago9</i>	<i>A. thaliana</i>	Mutation in gene involved in a small RNA pathway	Apospory like	Olmedo-Monfil et al., 2010
<i>rdr6</i>	<i>A. thaliana</i>	Mutation in gene involved in a small RNA pathway	Apospory like	Olmedo-Monfil et al., 2010
<i>sgs3</i>	<i>A. thaliana</i>	Mutation in gene involved in a small RNA pathway	Apospory like	Olmedo-Monfil et al., 2010
<i>mem</i>	<i>A. thaliana</i>	Mutation in gene encoding a RNA-helicase	Apospory like	Schmidt et al., 2011
<i>dmt102</i>	<i>Z. mays</i>	Mutation in gene involved in DNA methylation	Apospory like	Garcia-Aguilar et al., 2010
<i>dmt103</i>	<i>Z. mays</i>	Mutation in gene involved in DNA methylation	Apospory like	Garcia-Aguilar et al., 2010
<i>ago104</i>	<i>Z. mays</i>	Mutation in gene involved in a small RNA pathway	Diplospory like	Singh et al., 2011

A. thaliana, *Arabidopsis thaliana*; *Z. mays*, *Zea mays*.

Box 2. The evolution of apomixis and meiosis

Evidence suggest that apomixis evolved from a deregulation of the sexual pathway several times independently (Koltunow, 1993; Vielle-Calzada et al., 1996; Leblanc et al., 1997; Grimanelli et al., 2001; Grossniklaus, 2001; Tucker et al., 2003; Koltunow and Grossniklaus, 2003; Sharbel et al., 2009, 2010). Deregulation of genetic and epigenetic regulatory pathways has been hypothesized to be a consequence of hybridization and polyploidization, which have been proposed as preconditions for apomixis to occur (Asker and Jerling, 1992; Grossniklaus, 2001; Spillane et al., 2001; de Storme and Geelen, 2013).

Interestingly, according to a recent hypothesis, meiosis as a precondition for sexual reproduction is thought to have evolved as a repair mechanism for DNA damage induced by oxidative stress and ROS, and it has been proposed that the redox chemistry between oxidized DNA and the meiotic protein SPO11 is required for the generation of double-strand breaks, which are required for meiotic recombination and the repair of damaged DNA (Hörandl and Hadacek, 2013). However, low levels of ROS promote the acquisition of meiotic fate in maize anthers (Kelliher and Walbot, 2012). In line with this hypothesis, the metabolism of polyamine and spermidine, which are quenchers of ROS activity, is enriched in the AIC in *B. gunnisoniana* (Schmidt et al., 2014). Similarly, increasing evidence suggests the importance of the redox state for the development of the anther and male germline (reviewed by Zhang and Yang, 2014). Nevertheless, the role of stress and reactive oxygen species on regulating sexual versus apomictic reproduction remains unknown.

Increasing evidence highlights the importance of *AGO* activity in plant germline development and gamete formation (Nonomura et al., 2007; Wüest et al., 2010; Olmedo-Monfil et al., 2010; Singh et al., 2011; Borges et al., 2011; Tucker et al., 2012). This is reminiscent of the role of AGO proteins in the animal germline; proteins of the animal-specific PIWI-clade protect the genomic integrity of the germline, in particular by repressing the activity of transposons in invertebrates, although their role in vertebrates is less clear (Clark and Lau, 2014). AGO/PIWI proteins, and potentially other proteins involved in small RNA pathways, interact with VASA or VASA-like RNA helicases that are preferentially expressed in the germline (Yajima and Wessel, 2011). Although neither the PIWI clade of AGOs nor VASA RNA helicases have been identified in plants, recent evidence suggests that similar regulatory mechanisms evolved in the plant reproductive lineage, likely by convergence (Wüest et al., 2010; Schmidt et al., 2011).

In *Arabidopsis*, a role for *AGO9* and *MNEME (MEM)*, a RNA helicase that is preferentially expressed in the MMC, during germline specification has recently been described (Fig. 3; Table 2; Olmedo-Monfil et al., 2010; Schmidt et al., 2011). In plants heterozygous for *mem-1* or *mem-2*, more than one subepidermal enlarged cell instead of the single MMC develops in ~21% of ovules, whereas 37–48% of the ovules in *ago9* homozygotes show a similar phenotype, depending on the allele (Olmedo-Monfil et al., 2010; Schmidt et al., 2011). In *mem* and *ago9* mutants, these additional subepidermal enlarged cells directly give rise to a female gametophyte in the absence meiosis, closely resembling apospory (Olmedo-Monfil et al., 2010; Schmidt et al., 2011). Although the AGO9 protein has been detected only in the L1 layer of the developing ovule, and not in the MMC, *MEM* transcripts are highly enriched in the MMC but are also detected in the surrounding ovule tissue, albeit at much lower levels (Fig. 3; Olmedo-Monfil et al., 2010; Schmidt et al., 2011). Thus, *MEM* and *AGO9* likely repress the acquisition of reproductive fate in the surrounding tissues in a non-cell-autonomous manner (Olmedo-Monfil et al., 2010; Schmidt et al., 2011). Interestingly, *AGO9* plays a role in

repressing transposons in the germline, reminiscent of the role played by PIWI proteins in animals (Olmedo-Monfil et al., 2010). It remains to be determined whether *MEM* – like *VASA* in the animal germline – is involved in this process, potentially acting by aiding the unwinding of RNAs prior to their association with AGO proteins.

Similar phenotypes have also been reported for mutations that disrupt *RNA-DEPENDENT RNA POLYMERASE6 (RDR6)* and *SUPPRESSOR OF GENE SILENCING3 (SGS3)*, which are known to be required for the biogenesis of *trans*-acting siRNAs (Table 2; Olmedo-Monfil et al., 2010). In addition, features of apospory have been reported for the maize *dmt102* and *dmt103* lines, which carry mutations in the homologues of the *Arabidopsis* *CHROMOMETHYLTRANSFERASE3 (CMT3)* and *DOMAINS REARRANGED METHYLASE1 (DRM1)* and *DRM2*, respectively (Table 2; Garcia-Aguilar et al., 2010). Currently, no evidence has been reported that demonstrates the formation of viable offspring from the unreduced, supernumerous gametophytes seen in the maize *dmt102* and *dmt103* mutants, or in the *Arabidopsis* *ago9* and *mem* mutants. By contrast, a mutation in maize *AGO104*, a homologue of *Arabidopsis* *AGO9*, leads to features of diplospory and to the formation of triploid and tetraploid offspring following the fertilization of unreduced gametophytes (Table 2; Singh et al., 2011). The *ago104* phenotype is caused by a mutation that leads to defects in chromosome condensation during meiosis (affecting mega- and microsporogenesis) and, subsequently, to the formation




Fig. 3. The expression patterns of proteins/genes whose perturbations mimic apomeiosis. The expression patterns or abundance of protein are schematically shown for: (A) *MEL1* (Nonomura et al., 2007); (B) *MEM* (Schmidt et al., 2011); (C) *AGO5* (Tucker et al., 2012) and *AGO104* (Singh et al., 2011); and (D) *AGO9* (Olmedo-Monfil et al., 2010). During female germline formation, *MEL1* is expressed in the MMC, suggesting a cell-autonomous effect to cause failure of meiosis (Nonomura et al., 2007). However, a more complex regulation cannot be excluded, as in rare cases germline formation fails. *AGO104* and *AGO5* are both localized in the nucellus tissue, suggesting signalling from the sporophytic tissues of the nucellus to the developing germline (Singh et al., 2011; Tucker et al., 2012). Interestingly, *AGO9* was described to be restricted to the L1 layer of the nucellus, suggesting a non-cell-autonomous mechanism (Olmedo-Monfil et al., 2010). By contrast, highest expression of *MEM* has been observed in the MMC, so that a non-cell-autonomous mechanism to repress germline fate in the surrounding cells is likely. However, as *MEM* is also expressed in the nucellus surrounding the MMC at low levels, other mechanisms cannot be excluded.

of unreduced dyads (Singh et al., 2011). Interestingly, *AGO104* is expressed not in the MMC but in the surrounding somatic tissues, suggesting that the meiotic defect is mediated by a mobile signal (Fig. 3; Singh et al., 2011). Effects of AGO activity on meiosis have been described previously: in rice, mutations in the MMC-expressed gene *MEIOSIS ARRESTED AT LEPTOTENE1 (MEL1)* (Fig. 3) lead to meiotic arrest and sterility (affecting mega- and microsporogenesis) (Nonomura et al., 2007). MEL1 is closely related only to *Arabidopsis* AGO5 and thus belongs to a different AGO clade than AGO104 (Nonomura et al., 2007). Together, the data suggest diverse and important functions for AGO proteins that are active in different small RNA-dependent regulatory pathways during germline specification and meiosis in different plant species.

Predominant expression of *AGO1*, *AGO2*, *AGO5*, *AGO8* and *AGO9* has also been observed in the *Arabidopsis* egg cell (Wüest et al., 2010), although the function of these AGO proteins in the egg cell remains to be elucidated. *AGO5* is also highly enriched in sperm (Borges et al., 2011), and a role for AGO5 in a putative miRNA complex in the male germline has been proposed (Borges et al., 2011). During megasporogenesis, *AGO5* can be detected in sporophytic ovule tissues, but not in the developing female germline, similar to *AGO9* (Fig. 3; Tucker et al., 2012). Furthermore, plants carrying the semi-dominant *ago5-4* allele do not initiate female gametophyte development, suggesting that this particular mutation inhibits a somatic small RNA pathway that promotes the initiation of gametogenesis (Tucker et al., 2012).

In addition to involving DNA methylation and small RNA pathways, epigenetic pathways regulate chromatin organization and histone modifications. It was recently reported that large-scale chromatin reprogramming establishes an epigenetic and transcriptional state in the *Arabidopsis* MMC that is distinct from that in the surrounding tissue (She et al., 2013). These changes likely contribute to the acquisition of germline fate and to the transition to the gametophytic phase, rather than being only a precondition for meiosis (She et al., 2013). This is supported by the finding of similar histone modifications and histone variant dynamics in the additional subepidermal enlarged cells in *ago9*, *sds3* and *rdr6* mutants (She et al., 2013).

In conclusion, epigenetic regulatory pathways play important roles during the acquisition of germline fate, during germline differentiation and for discriminating between a meiotic and a mitotic fate. It remains unknown whether the influence of stress on germline specification described above acts via changes in the activity of epigenetic pathways or through independent mechanisms.

Polarity and cell fate determination during megagametogenesis

Whether generated by sexual reproduction or apomixis, a highly polarized structure harbouring functionally distinct cell types is established during megasporogenesis from a single FMS by only two or three mitotic divisions and cellularization. Recent studies in *Arabidopsis* have identified a number of factors that can influence this polarity and the subsequent development of the gametophyte (summarized in Table 3; reviewed by Sundaresan and Alandete-Saez, 2010; Sprunck and Gross-Hardt, 2011; Lituiev and Grossniklaus, 2014).

Factors regulating FMS selection

To initiate megagametogenesis, typically only the chalazal-most spore in the tetrad survives and differentiates into the FMS, but the mechanism governing FMS selection and survival is unclear. The *Arabidopsis antikeyvorkian* mutant affects FMS selection but the

corresponding gene remains to be cloned (Yang and Sundaresan, 2000). More recently, *ARABINOGLACTAN PROTEIN18 (AGP18)* was found to be important for the survival and selection of the FMS (Table 3; Acosta-García and Vielle-Calzada, 2004; Demesa-Arévalo and Vielle-Calzada, 2013). Overexpression of *AGP18* in ovules results in the survival of more than one of the four megaspores (Demesa-Arévalo and Vielle-Calzada, 2013). The mechanism by which *AGP18* determines FMS selection remains unknown, although it has been hypothesized that AGP proteins, which are attached to the plasma membrane through a glycosylphosphatidylinositol (GPI) anchor, can act as components of signalling pathways (Youl et al., 1998; Börner et al., 2003; Ellis et al., 2010; Seifert and Roberts, 2007; Zhang et al., 2011).

Subsequent to megaspore selection, the FMS forms the mature gametophyte, which harbours four functionally distinct cell types, typically through three mitotic divisions. How cell fate acquisition is regulated and when cell fate is determined during this process is still largely unclear. It has been proposed that positional information might be involved in the determination of cell fate (Grossniklaus and Schneitz, 1998; Sundaresan and Alandete-Saez, 2010; Sprunck and Gross-Hardt, 2011; Lituiev and Grossniklaus, 2014). During the syncytial phase, nuclei migrate and occupy predefined positions in the female gametophyte. In mutants with supernumerary nuclei, the position of the nuclei along the micropylar-chalazal axis of the embryo sac affects their cell fate, indicating that they are influenced by positional information (Table 3; Gross-Hardt et al., 2007; Pagnussat et al., 2007; Moll et al., 2008a; Moll et al., 2008b; Johnston et al., 2010).

The role of auxin and cytokinin in establishing and maintaining polarity

It has been proposed that the plant hormone auxin plays a pivotal role in establishing and maintaining polarity by forming a gradient in the developing embryo sac (Pagnussat et al., 2009). The auxin gradient was thought to be mediated by auxin influx from sporophytic tissues at early stages and by localized biosynthesis at later stages of female gametophyte development. Abolishing the auxin gradient, by expressing the YUCCA1 (YUC1) auxin biosynthetic protein in the entire embryo sac or by modulating the auxin response by downregulating selected *AUXIN RESPONSE FACTOR (ARF)* genes, led to the loss or, at low frequencies, the mis-expression of cell fate markers in the female gametophyte (Table 3; Pagnussat et al., 2009). However, theoretical models attempting to describe the auxin gradient in the female gametophyte showed that only very shallow auxin gradients can be established even when using the most favourable parameters (Lituiev et al., 2013). A sensitivity analysis demonstrated that the steepness of the obtained gradients is not sufficient to determine distinct cell fates (Lituiev et al., 2013). Furthermore, the reinvestigation of auxin signalling using various sensors failed to detect auxin in the female gametophyte but instead found auxin signalling to be restricted to the surrounding ovule tissues in a dynamic polar pattern (Ceccato et al., 2013; Lituiev et al., 2013). This polar auxin pattern in sporophytic tissues may non-cell-autonomously influence cell specification in the female gametophyte and may have been affected by manipulating the expression of *YUC1* and *ARFs* (Lituiev et al., 2013).

Auxin signalling is interrelated with the cytokinin pathway (Müller and Sheen, 2008; Bencivenga et al., 2012; Cheng et al., 2013) and, not surprisingly therefore, cytokinin signalling has also been shown to play a role in germline development. For example, cytokinin levels influence ovule patterning by affecting the

Table 3. Mutations involved in polarity and cell fate determination in the *Arabidopsis thaliana* female gametophyte

Mutation	Description	Phenotype	Reference
<i>cki1</i>	Mutation in a gene causing cytokinin-independent activation of the cytokinin signalling pathway	Arrest starting from FG4	Pischke et al., 2002; Hejátko et al., 2003
<i>agp18</i>	RNAi targeting <i>ARABINOGALACTAN PROTEIN18</i> transcripts	Arrest at FG1	Acosta-García and Vielle-Calzada, 2004
<i>rbr1</i>	Mutation in a core cell cycle regulator gene	Nuclear overproliferation	Ebel et al., 2004; Johnston et al., 2010
<i>lis</i>	Mutation in a gene encoding a component of the RNA splicing machinery	Synergids and the central cell adopt an egg cell-like fate	Gross-Hardt et al., 2007; Völz et al., 2012
<i>eostre</i>	Mutation leading to the misexpression of <i>BLH1</i>	One synergid cell differentiates into an additional egg cell	Pagnussat et al., 2007
<i>clo/gfa1</i>	Mutation in a gene encoding a component of the RNA splicing machinery	Synergids and the central cell adopt an egg cell-like fate	Moll et al., 2008b
<i>ato</i>	Mutation in a gene encoding a component of the RNA splicing machinery	Synergids and the central cell adopt an egg cell-like fate	Moll et al., 2008b
<i>amiR-ARFa</i>	amiRNA targeting transcripts of the auxin signalling pathway	Synergid identity lost, partly adopting an egg cell-like fate	Pagnussat et al., 2009
<i>ahk2-7, ahk3-3, cre1-12</i>	Triple mutant in genes encoding components of the cytokinin signalling pathway	Arrest at FG1	Cheng et al., 2013
<i>ahp1, ahp2-1, ahp3, ahp4, ahp5</i>	Multiple mutant in genes encoding components of the cytokinin signalling pathway	Arrest at FG7	Cheng et al., 2013
<i>hda7</i>	Mutation in a histone deacetylase gene	Arrest at FG4	Cigliano et al., 2013
<i>myb64, myb119</i>	Double mutant in MYB transcription factor genes	Arrest during FG5 transition	Rabiger and Drews, 2013

amiRNA, artificial microRNA; *BLH1*, *BELL1-LIKE HOMEODOMAIN1*; RNAi, RNA interference. Female gametophyte stages (FG) are according to Christensen et al. (1997).

expression of *PIN-FORMED1* (*PIN1*), which encodes an auxin efflux carrier (Bencivenga et al., 2012; Luschnig and Vert, 2014). This is consistent with the finding that cytokinin regulates *PIN1* expression in roots (Dello Ioio et al., 2008; Ruzicka et al., 2009). In ovules, the regulatory pathway involves two transcription factors, *SPOROCTLESS/NOZZLE* (*SPL/NZZ*) and the homeodomain protein *BELL1* (*BEL1*). Mutations in the gene encoding *SPL/NZZ*, which is required for the initiation of megasporogenesis, lead to reduced expression of *PIN1*, while the effects of exogenous cytokinin are mediated by *BEL1*, which is important for ovule identity, and lead to an altered pattern of auxin signalling in the ovule (Table 3; Schiefthaler et al., 1999; Yang et al., 1999; Balasubramanian and Schneitz, 2000; Sieber et al., 2004; Brambilla et al., 2007; Bencivenga et al., 2012). In addition to *PIN1*, *PIN3* is expressed during ovule development (Ceccato et al., 2013). However, no effect on ovule or female germline development has been reported in *pin3* mutants and a potential functional interaction between cytokinin and *PIN3* has not been investigated (Ceccato et al., 2013). Thus, although an auxin gradient in the embryo sac could not be confirmed, auxin and cytokinin do play important roles in the sporophytic tissues of the ovule.

Consistent with the crosstalk between the auxin and cytokinin pathways, cytokinin is involved in communication between sporophytic ovule tissues and the developing female gametophyte (Table 3; Cheng et al., 2013). In *Arabidopsis*, different cytokinin receptors expressed in the chalazal ovule tissues act redundantly to regulate FMS specification (Table 3; Cheng et al., 2013). During megagametogenesis, the histidine protein kinase *CYTOKININ-INDEPENDENT1* (*CKI1*) has important functions, and *cki1* mutants affect the mitotic divisions during gametophyte development (Table 3; Pischke et al., 2002; Hejátko et al., 2003; Cheng et al., 2013). Although related to the *Arabidopsis* histidine kinases (AHKs) *AHK2*, *AHK3* and *AHK4*, which act as cytokinin receptors, *CKI1* lacks a cytokinin-binding domain and activates the cytokinin signalling pathway in the absence of cytokinin (Kakimoto, 1996; Nakamura et al., 1999; Urao et al., 2000; Hwang

and Sheen, 2001; Yamada et al., 2001; Mähönen et al., 2006). *Arabidopsis* double mutants affecting *MYB-DOMAIN PROTEIN64* (*MYB64*) and *MYB119*, which encode two closely related R2R3-MYB domain transcription factors, also display a *cki1*-like phenotype (Table 3; Rabiger and Drews, 2013). Double mutant *myb64 myb119* gametophytes undergo extra mitotic division cycles and usually fail to cellularize (Rabiger and Drews, 2013). In the few cellularized mutant embryo sacs, cell fate is not properly established and the polarity of the embryo sac is affected (Rabiger and Drews, 2013). Furthermore, while *MYB64* and *MYB119* act redundantly during female gametophyte development, *MYB119* but not *MYB64* is regulated by *CKI1* (Rabiger and Drews, 2013).

Epigenetic and post-transcriptional regulation of gametophyte development

In addition to hormonal pathways and transcription factors, epigenetic regulators are involved in establishing polarity in the developing gametophyte. Recently, a role for HISTONE DEACETYLASE7 (*HDA7*) during megagametogenesis and embryo development has been demonstrated (Cigliano et al., 2013). In *hda7-2* mutants at the four-nucleate stage of megagametogenesis, the two nuclei located at the micropylar pole degenerate, suggesting that histone deacetylation is required for survival and possibly for fate determination of the micropylar nuclei (Table 3; Cigliano et al., 2013). Other important gene regulatory control mechanisms involve the storage of mRNAs in mRNA-protein complexes, mRNA processing and mRNA degradation (reviewed by Hafidh et al., 2011). Regulation of the asymmetric distribution and processing of mRNAs involving RNA-binding proteins is known to be a determinant of protein gradients, cell polarity, cell fate decisions and patterning during development (Hafidh et al., 2011). For example, this is well described in *Drosophila* embryo genesis but also relevant for polar pollen tube growth in plants (Hafidh et al., 2011). In agreement with the emerging roles of mRNA storage and processing, components of the RNA splicing machinery have been identified as being crucial for cell type specification and the restriction

of gametic fate in the *Arabidopsis* embryo sac (Table 3; Gross-Hardt et al., 2007; Moll et al., 2008b; Völz et al., 2012). In *lachesis* (*lis*) mutants, the expression of a marker for egg cell identity extends to adjacent gametophytic cells, the synergids and the central cell (Table 3; Gross-Hardt et al., 2007). As the phenotype becomes stronger as time progresses, LIS may predominantly play a role in maintaining egg cell identity. LIS encodes a homologue of the yeast splicing factor PRP4 (Gross-Hardt et al., 2007). Similar to LIS, CLOTHO/GAMETOPHYTIC FACTOR1 (CLO/GFA1) and ATROPUS (ATO) are also important for restricting gametic fate in the mature gametophyte (Table 3; Moll et al., 2008b). CLO/GFA1 encodes a homologue of Sn114, an essential component of the spliceosome, while ATO encodes the *Arabidopsis* homologue of SF3a60, which plays a role in pre-spliceosome formation (Moll et al., 2008b). The activities of LIS and CLO are related, as CLO is important for the tissue specificity of LIS expression (Moll et al., 2008b). LIS is strongly enriched in female gametes, suggesting that it regulates the maintenance of cell fate by lateral inhibition of the adjacent accessory cells in the female gametophyte, the synergid and antipodal cells (Gross-Hardt et al., 2007; Moll et al., 2008b; Völz et al., 2012). In summary, the splicing machinery is important for the specification and maintenance of cellular identity in the female gametophyte. Whether this is mediated through specific effects of some of its components in the embryo sac or caused by a general deficiency in splicing – also affecting pre-mRNAs of cell specification factors – remains to be determined.

In conclusion, although many mutants that exhibit disrupted embryo sac polarity or cell type-specific expression have been identified over the past decade, we are still far from understanding these processes at the molecular level. Currently, we have a partial list of components involved in cell specification but we do not understand how they work together to pattern the female gametophyte. Importantly, many of the observed effects may be indirect, e.g. caused by the mis-positioning of nuclei in the embryo sac, or the identified factors act after the initial specification of cell fate, in the maintenance of cell identity or during cell differentiation. A clear candidate for a cell fate determinant – one that cell-autonomously specifies cell type identity – is still being sought after. Transcription factors of the RKD family can at least partially reprogramme sporophytic cells towards an egg cell fate when overexpressed (Koszegi et al., 2011) but, owing to genetic redundancy, functional analyses of these transcription factors proved difficult and their potential gametophytic phenotypes are unknown.

Conclusions

The male and female plant germlines are ideal models for studying the role of polarity, cell specification processes and the transition from sporophytic to gametophytic fate, which is a key step in the plant life cycle. Apart from being scientifically fascinating, understanding the molecular mechanisms underlying the specification and development of plant reproductive lineages is relevant for targeted manipulations of reproduction for crop improvement and seed production, in particular to achieve the longstanding goal of engineering apomixis in crop plants. Important aspects will be to determine whether distinct or similar genetic and epigenetic modifications govern apomixis in different species, involving aposporous and diplosporous accessions, and to identify common features.

Recent studies have yielded important insights into various aspects of sexual and apomictic germline formation, providing a glimpse of the complex regulatory mechanisms required to control reproductive development in plants. In this Review, we have discussed studies that address the genetic basis underlying the transition from somatic to

germline fate, the repression of additional reproductive lineages, and cell specification during megagametogenesis, which together ensure reproductive success. As highlighted above, many different pathways are involved, including hormonal pathways, epigenetic regulation via small RNAs, transcriptional regulation by transcription factors and post-transcriptional control mechanisms. However, we currently do not know how these pathways are interconnected to form regulatory networks. Building on recent investigations, an important aim of future research will be to compare whole transcriptome analyses and combine these with detailed studies of the molecular mechanisms at play in different species, while taking evolutionary aspects into consideration.

Finally, little is known about the influence of stress or changing environmental conditions on germline specification. Interestingly, heat stress as well as the abundance of ROS were reported to influence the meiotic versus apomeiotic fate decision of spore mother cells. However, contradicting hypotheses have been portrayed, highlighting a need for more investigations in this area. In the longer term, this may be important not only for a better understanding of the regulatory networks underlying reproductive development and their interactions with environmental factors, but also for the improvement of agricultural plants under changing climate conditions.

Acknowledgements

We thank our colleagues in the Grossniklaus laboratory for interesting discussions and three anonymous reviewers for their helpful comments, which helped us to improve the manuscript.

Competing interests

The authors declare no competing financial interests.

Funding

Work on gametophyte development, apomixis and epigenetic gene regulation in U.G.'s laboratory is supported by the University of Zürich, and by grants from the 'Staatssekretariat für Bildung und Forschung' in the framework of COST action FA0903 (to U.G. and A.S.), the Swiss National Science Foundation (to U.G.) and the European Research Council (to U.G.).

References

- Abiko, M., Maeda, H., Tamura, K., Hara-Nishimura, I. and Okamoto, T. (2013). Gene expression profiles in rice gametes and zygotes: identification of gamete-enriched genes and up- or down-regulated genes in zygotes after fertilization. *J. Exp. Bot.* **64**, 1927–1940.
- Acosta-García, G. and Vielle-Calzada, J.-P. (2004). A classical arabinogalactan protein is essential for the initiation of female gametogenesis in *Arabidopsis*. *Plant Cell* **16**, 2614–2628.
- Albrecht, C., Russinova, E., Hecht, V., Baaijens, E. and de Vries, S. (2005). The *Arabidopsis thaliana* SOMATIC EMBRYOGENESIS RECEPTOR-LIKE KINASES 1 and 2 control male sporogenesis. *Plant Cell* **17**, 3337–3349.
- Anderson, S. N., Johnson, C. S., Jones, D. S., Conrad, L. J., Gou, X., Russell, S. D. and Sundaresan, V. (2013). Transcriptomes of isolated *Oryza sativa* gametes characterized by deep sequencing: evidence for distinct sex-dependent chromatin and epigenetic states before fertilization. *Plant J.* **76**, 729–741.
- Asker, S. and Jerling, E. L. (1992). *Apomixis in Plants*. Boca Raton, Florida, USA: CRC Press.
- Bachelier, J. B. and Friedman, W. E. (2011). Female gamete competition in an ancient angiosperm lineage. *Proc. Natl. Acad. Sci. USA* **108**, 12360–12365.
- Balasubramanian, S. and Schneitz, K. (2000). NOZZLE regulates proximal-distal pattern formation, cell proliferation and early sporogenesis during ovule development in *Arabidopsis thaliana*. *Development* **127**, 4227–4238.
- Bencivenga, S., Simonini, S., Benková, E. and Colombo, L. (2012). The transcription factors BEL1 and SPL are required for cytokinin and auxin signaling during ovule development in *Arabidopsis*. *Plant Cell* **24**, 2886–2897.
- Berger, F. and Twell, D. (2011). Germline specification and function in plants. *Annu. Rev. Plant Biol.* **62**, 461–484.
- Bicknell, R. A. and Koltunow, A. M. (2004). Understanding apomixis: recent advances and remaining conundrums. *Plant Cell* **16** Suppl. 1, S228–S245.
- Boavida, L. C., Becker, J. D. and Feijo, J. A. (2005). The making of gametes in higher plants. *Int. J. Dev. Biol.* **49**, 595–614.
- Borg, M., Brownfield, L. and Twell, D. (2009). Male gametophyte development: a molecular perspective. *J. Exp. Bot.* **60**, 1465–1478.

- Borges, F., Pereira, P. A., Slotkin, R. K., Martienssen, R. A. and Becker, J. D. (2011). MicroRNA activity in the *Arabidopsis* male germline. *J. Exp. Bot.* **62**, 1611–1620.
- Borner, G. H. H., Lilley, K. S., Stevens, T. J. and Dupree, P. (2003). Identification of glycosylphosphatidylinositol-anchored proteins in *Arabidopsis*. A proteomic and genomic analysis. *Plant Physiol.* **132**, 568–577.
- Brambilla, V., Battaglia, R., Colombo, M., Masiero, S., Bencivenga, S., Kater, M. M. and Colombo, L. (2007). Genetic and molecular interactions between BELL1 and MADS box factors support ovule development in *Arabidopsis*. *Plant Cell* **19**, 2544–2556.
- Brownfield, L. and Köhler, C. (2011). Unreduced gamete formation in plants: mechanisms and prospects. *J. Exp. Bot.* **62**, 1659–1668.
- Canales, C., Bhatt, A. M., Scott, R. and Dickinson, H. (2002). EXS, a putative LRR receptor kinase, regulates male germline cell number and tapetal identity and promotes seed development in *Arabidopsis*. *Curr. Biol.* **12**, 1718–1727.
- Ceccato, L., Masiero, S., Sinha Roy, D., Bencivenga, S., Roig-Villanova, I., Dittengou, F. A., Palme, K., Simon, R. and Colombo, L. (2013). Maternal control of PIN1 is required for female gametophyte development in *Arabidopsis*. *PLoS ONE* **8**, e66148.
- Cheng, C.-Y., Mathews, D. E., Schaller, G. E. and Kieber, J. J. (2013). Cytokinin-dependent specification of the functional megaspore in the *Arabidopsis* female gametophyte. *Plant J.* **73**, 929–940.
- Chettoor, A. M., Givan, S. A., Cole, R. A., Coker, C. T., Unger-Wallace, E., Vejlupekova, Z., Vollbrecht, E., Fowler, J. E. and Evans, M. M. S. (2014). Discovery of novel transcripts and gametophytic functions via RNA-seq analysis of maize gametophytic transcriptomes. *Genome Biol.* **15**, 414.
- Christensen, C. A., King, E. J., Jordan, J. R. and Drews, G. N. (1997). Megagametogenesis in *Arabidopsis* wild type and the *Gf* mutant. *Sex. Plant Reprod.* **10**, 49–64.
- Cigliano, R. A., Cremona, G., Paparo, R., Termolino, P., Perrella, G., Gutzat, R., Consiglio, M. F. and Conicella, C. (2013). Histone deacetylase AtHDA7 is required for female gametophyte and embryo development in *Arabidopsis*. *Plant Physiol.* **163**, 431–440.
- Clark, J. P. and Lau, N. C. (2014). Piwi proteins and piRNAs step onto the systems biology stage. *Adv. Exp. Med. Biol.* **825**, 159–197.
- Colombet, J., Boisson-Dernier, A., Ros-Palau, R., Vera, C. E. and Schroeder, J. I. (2005). *Arabidopsis* SOMATIC EMBRYOGENESIS RECEPTOR KINASES1 and 2 are essential for tapetum development and microspore maturation. *Plant Cell* **17**, 3350–3361.
- Crismani, W., Girard, C. and Mercier, R. (2013). Tinkering with meiosis. *J. Exp. Bot.* **64**, 55–65.
- De Storme, N. and Geelen, D. (2013). Sexual polyploidization in plants - cytological mechanisms and molecular regulation. *New Phytol.* **198**, 670–684.
- De Storme, N. and Geelen, D. (2014). The impact of environmental stress on male reproductive development in plants: biological processes and molecular mechanisms. *Plant Cell Environ.* **37**, 1–18.
- Dello Ioio, R., Nakamura, K., Moubayidin, L., Perilli, S., Taniguchi, M., Morita, M. T., Aoyama, T., Costantino, P. and Sabatini, S. (2008). A genetic framework for the control of cell division and differentiation in the root meristem. *Science* **322**, 1380–1384.
- Demesa-Arévalo, E. and Velle-Calzada, J.-P. (2013). The classical arabinogalactan protein AGP18 mediates megaspore selection in *Arabidopsis*. *Plant Cell* **25**, 1274–1287.
- Drews, G. N. and Koltunow, A. M. G. (2011). The female gametophyte. *Arabidopsis Book* **9**, e0155.
- d'Erfurth, I., Jolivet, S., Froger, N., Catrice, O., Novatchkova, M. and Mercier, R. (2009). Turning meiosis into mitosis. *PLoS Biol.* **7**, e1000124.
- d'Erfurth, I., Cromer, L., Jolivet, S., Girard, C., Horlow, C., Sun, Y., To, J. P. C., Berchowitz, L. E., Copenhaver, G. P. and Mercier, R. (2010). The cyclin-A CYCA1;2/TAM is required for the meiosis I to meiosis II transition and cooperates with OSD1 for the prophase to first meiotic division transition. *PLoS Genet.* **6**, e1000989.
- Ebel, C., Mariconti, L. and Gruissem, W. (2004). Plant retinoblastoma homologues control nuclear proliferation in the female gametophyte. *Nature* **429**, 776–780.
- Ellis, M., Egelund, J., Schultz, C. J. and Bacic, A. (2010). Arabinogalactan-proteins: key regulators at the cell surface? *Plant Physiol.* **153**, 403–419.
- Feng, X. and Dickinson, H. G. (2010). Tapetal cell fate, lineage and proliferation in the *Arabidopsis* anther. *Development* **137**, 2409–2416.
- García-Aguilar, M., Michaud, C., Leblanc, O. and Grimanelli, D. (2010). Inactivation of a DNA methylation pathway in maize reproductive organs results in apomixis-like phenotypes. *Plant Cell* **22**, 3249–3267.
- Germanà, M. A. (2011). Gametic embryogenesis and haploid technology as valuable support to plant breeding. *Plant Cell Rep.* **30**, 839–857.
- Grimanelli, D., Leblanc, O., Perotti, E. and Grossniklaus, U. (2001). Developmental genetics of gametophytic apomixis. *Trends Genet.* **17**, 597–604.
- Gross-Hardt, R., Kägi, C., Baumann, N., Moore, J. M., Baskar, R., Gagliano, W. B., Jürgens, G. and Grossniklaus, U. (2007). LACHESIS restricts gametic cell fate in the female gametophyte of *Arabidopsis*. *PLoS Biol.* **5**, e47.
- Grossniklaus, U. (2001). From sexuality to apomixis: molecular and genetic approaches. In *The Flowering of Apomixis: From Mechanisms to Genetic Engineering* (ed. Y. Savidan, J. Carman and T. Dresselhaus), pp. 168–211. Mexico, DF: CIMMYT.
- Grossniklaus, U. (2011). Plant germline development: a tale of cross-talk, signaling, and cellular interactions. *Sex. Plant Reprod.* **24**, 91–95.
- Grossniklaus, U. and Schneitz, K. (1998). The molecular and genetic basis of ovule and megagametophyte development. *Semin. Cell Dev. Biol.* **9**, 227–238.
- Grossniklaus, U., Koltunow, A. and van Lookeren Campagne, M. (1998a). A bright future for apomixis. *Trends Plant Sci.* **3**, 415–416.
- Grossniklaus, U., Moore, J. M. and Gagliano, W. B. (1998b). Molecular and genetic approaches to understanding and engineering apomixis: *Arabidopsis* as a powerful tool. In *Advances in Hybrid Rice Technology. Proceedings of the 3rd International Symposium on Hybrid Rice 1996* (ed. S. S. Virmani, E. A. Siddiq and K. Muralidharan), pp. 187–211. Manila, Philippines: International Rice Research Institute.
- Gutierrez-Marcos, J. F. and Dickinson, H. G. (2012). Epigenetic reprogramming in plant reproductive lineages. *Plant Cell Physiol.* **53**, 817–823.
- Hafidh, S., Čapková, V. and Honys, D. (2011). Safe keeping the message: mRNP complexes tweaking after transcription. *Adv. Exp. Med. Biol.* **722**, 118–136.
- Haig, D. (1990). New perspectives on the angiosperm female gametophyte. *Botanical Rev.* **56**, 236–274.
- Hejác, J., Pernisová, M., Eneva, T., Palme, K. and Brzobohatý, B. (2003). The putative sensor histidine kinase CK1I is involved in female gametophyte development in *Arabidopsis*. *Mol. Genet. Genomics* **269**, 443–453.
- Hörandl, E. and Hadacek, F. (2013). The oxidative damage initiation hypothesis for meiosis. *Plant Reprod.* **26**, 351–367.
- Huang, B.-Q. and Russell, S. D. (1992). Female germ unit: organization, isolation, and function. *Int. Rev. Cytol.* **140**, 233–293.
- Huang, B. Q. and Sheridan, W. F. (1996). Embryo sac development in the maize *indeterminate gametophyte1* mutant: abnormal nuclear behavior and defective microtubule organization. *Plant Cell* **8**, 1391–1407.
- Hwang, I. and Sheen, J. (2001). Two-component circuitry in *Arabidopsis* cytokinin signal transduction. *Nature* **413**, 383–389.
- Jia, G., Liu, X., Owen, H. A. and Zhao, D. (2008). Signaling of cell fate determination by the TPD1 small protein and EMS1 receptor kinase. *Proc. Natl. Acad. Sci. USA* **105**, 2220–2225.
- Johnston, A. J., Kirioukhova, O., Barrell, P. J., Rutten, T., Moore, J. M., Baskar, R., Grossniklaus, U. and Gruissem, W. (2010). Dosage-sensitive function of *RETINOBLASTOMA RELATED* and convergent epigenetic control are required during the *Arabidopsis* life cycle. *PLoS Genet.* **6**, pe1000988.
- Kakimoto, T. (1996). CK1I, a histidine kinase homolog implicated in cytokinin signal transduction. *Science* **274**, 982–985.
- Kelliher, T. and Walbot, V. (2012). Hypoxia triggers meiotic fate acquisition in maize. *Science* **337**, 345–348.
- Kelliher, T. and Walbot, V. (2014). Maize germinal cell initials accommodate hypoxia and precociously express meiotic genes. *Plant J.* **77**, 639–652.
- Koltunow, A. M. (1993). Apomixis: embryo sacs and embryos formed without meiosis or fertilization in ovules. *Plant Cell* **5**, 1425–1437.
- Koltunow, A. and Grossniklaus, U. (2003). Apomixis: a developmental perspective. *Annu. Rev. Plant Biol.* **54**, 547–574.
- Koltunow, A. M., Bicknell, R. A. and Chaudhury, A. M. (1995). Apomixis: molecular strategies for the generation of genetically identical seeds without fertilization. *Plant Physiol.* **108**, 1345–1352.
- Koltunow, A. M., Johnson, S. D., Rodrigues, J. C. M., Okada, T., Hu, Y., Tsuchiya, T., Wilson, S., Fletcher, P., Ito, K., Suzuki, G. et al. (2011). Sexual reproduction is the default mode in apomictic *Hieracium* subgenus *Pilosella*, in which two dominant loci function to enable apomixis. *Plant J.* **66**, 890–902.
- Kozegi, D., Johnston, A. J., Rutten, T., Czihal, A., Altschmied, L., Kumlehn, J., Wüst, S. E. J., Kirioukhova, O., Gheyselinck, J., Grossniklaus, U. et al. (2011). Members of the RKD transcription factor family induce an egg cell-like gene expression program. *Plant J.* **67**, 280–291.
- Leblanc, O., Armstead, I., Pessino, S., Ortiz, J. P. A., Evans, C., doValle, C. and Hayward, M. D. (1997). Non-radioactive mRNA fingerprinting to visualise gene expression in mature ovaries of *Bracharia* hybrids derived from *B. brizantha*, an apomictic tropical forage. *Plant Sci.* **126**, 49–58.
- Lin, B.-Y. (1984). Ploidy barrier to endosperm development in maize. *Genetics* **107**, 103–115.
- Lituiev, D. S. and Grossniklaus, U. (2014). Patterning of the angiosperm female gametophyte through the prism of theoretical paradigms. *Biochem. Soc. Trans.* **42**, 332–339.
- Lituiev, D. S., Krohn, N. G., Müller, B., Jackson, D., Hellriegel, B., Dresselhaus, T. and Grossniklaus, U. (2013). Theoretical and experimental evidence indicates that there is no detectable auxin gradient in the angiosperm female gametophyte. *Development* **140**, 4544–4553.
- Luschnig, C. and Vert, G. (2014). The dynamics of plant plasma membrane proteins: PINs and beyond. *Development* **141**, 2924–2938.
- Maheshwari, P. (1950). *An Introduction to the Embryology of Angiosperms*. New York: McGraw-Hill Publications.

- Mähönen, A. P., Higuchi, M., Törökangas, K., Miyawaki, K., Pischke, M. S., Sussman, M. R., Helariutta, Y. and Kakimoto, T. (2006). Cytokinins regulate a bidirectional phosphorelay network in *Arabidopsis*. *Curr. Biol.* **16**, 1116–1122.
- Mallory, A. and Vaucheret, H. (2010). Form, function, and regulation of ARGONAUTE proteins. *Plant Cell* **22**, 3879–3889.
- Marimuthu, M. P. A., Jolivet, S., Ravi, M., Pereira, L., Davda, J. N., Cromer, L., Wang, L., Nogué, F., Chan, S. W. L., Siddiqi, I. et al. (2011). Synthetic clonal reproduction through seeds. *Science* **331**, 876.
- Meister, G. (2013). Argonaute proteins: functional insights and emerging roles. *Nat. Rev. Genet.* **14**, 447–459.
- Moll, C., Nielsen, N. and Gross-Hardt, R. (2008a). Mutants with aberrant numbers of gametic cells shed new light on old questions. *Plant Biol. (Stuttg)* **10**, 529–533.
- Moll, C., von Lyncker, L., Zimmermann, S., Kägi, C., Baumann, N., Twill, D., Grossniklaus, U. and Gross-Hardt, R. (2008b). *CLO/GFA1* and *ATO* are novel regulators of gametic cell fate in plants. *Plant J.* **56**, 913–921.
- Müller, B. and Sheen, J. (2008). Cytokinin and auxin interaction in root stem-cell specification during early embryogenesis. *Nature* **453**, 1094–1097.
- Nakamura, A., Kakimoto, T., Imamura, A., Suzuki, T., Ueguchi, C. and Mizuno, T. (1999). Biochemical characterization of a putative cytokinin-responsive Histidine Kinase, CK11, from *Arabidopsis thaliana*. *Biosci. Biotechnol. Biochem.* **63**, 1627–1630.
- Nonomura, K.-I., Miyoshi, K., Eiguchi, M., Suzuki, T., Miyao, A., Hirochika, H. and Kurata, N. (2003). The *MSP1* gene is necessary to restrict the number of cells entering into male and female sporogenesis and to initiate anther wall formation in rice. *Plant Cell* **15**, 1728–1739.
- Nonomura, K.-I., Morohoshi, A., Nakano, M., Eiguchi, M., Miyao, A., Hirochika, H. and Kurata, N. (2007). A germ cell specific gene of the ARGONAUTE family is essential for the progression of premeiotic mitosis and meiosis during sporogenesis in rice. *Plant Cell* **19**, 2583–2594.
- Okada, T., Hu, Y., Tucker, M. R., Taylor, J. M., Johnson, S. D., Spriggs, A., Tsuchiya, T., Oelkers, K., Rodrigues, J. C. M. and Koltunow, A. M. (2013). Enlarging cells initiating apomixis in *Hieracium praealtum* transition to an embryo sac program prior to entering mitosis. *Plant Physiol.* **163**, 216–231.
- Olmedo-Monfil, V., Durán-Figueroa, N., Arteaga-Vázquez, M., Demesa-Arévalo, E., Autran, D., Grimanelli, D., Slotkin, R. K., Martienssen, R. A. and Vienne-Calzada, J.-P. (2010). Control of female gamete formation by a small RNA pathway in *Arabidopsis*. *Nature* **464**, 628–632.
- Pagnussat, G. C., Yu, H.-J. and Sundaresan, V. (2007). Cell-fate switch of synergist to egg cell in *Arabidopsis eostre* mutant embryo sacs arises from misexpression of the BEL1-like homeodomain gene *BLH1*. *Plant Cell* **19**, 3578–3592.
- Pagnussat, G. C., Alandete-Saez, M., Bowman, J. L. and Sundaresan, V. (2009). Auxin-dependent patterning and gamete specification in the *Arabidopsis* female gametophyte. *Science* **324**, 1684–1689.
- Pecix, Y., Rallo, G., Folzer, H., Cigna, M., Gudin, S. and Le Bris, M. (2011). Polyploidization mechanisms: temperature environment can induce diploid gamete formation in *Rosa* sp. *J. Exp. Bot.* **62**, 3587–3597.
- Pischke, M. S., Jones, L. G., Otsuga, D., Fernandez, D. E., Drews, G. N. and Sussman, M. R. (2002). An *Arabidopsis* histidine kinase is essential for megagametogenesis. *Proc. Natl. Acad. Sci. USA* **99**, 15800–15805.
- Rabiger, D. S. and Drews, G. N. (2013). MYB64 and MYB119 are required for cellularization and differentiation during female gametogenesis in *Arabidopsis thaliana*. *PLoS Genet.* **9**, e1003783.
- Raghavan, V. (1997). *Molecular Embryology of Flowering Plants*. Cambridge, UK: Cambridge University Press.
- Ravi, M., Marimuthu, M. P. A. and Siddiqi, I. (2008). Gamete formation without meiosis in *Arabidopsis*. *Nature* **451**, 1121–1124.
- Russell, S. D. (1979). Fine structure of megagametophyte development in *Zea mays*. *Can. J. Bot.* **57**, 1093–1110.
- Ruzicka, K., Simásková, M., Duclercq, J., Petrásek, J., Zázimalová, E., Simon, S., Friml, J., Van Montagu, M. C. E. and Benková, E. (2009). Cytokinin regulates root meristem activity via modulation of the polar auxin transport. *Proc. Natl. Acad. Sci. USA* **106**, 4284–4289.
- Schiefthaler, U., Balasubramanian, S., Sieber, P., Chevalier, D., Wisman, E. and Schneitz, K. (1999). Molecular analysis of *NOZZLE*, a gene involved in pattern formation and early sporogenesis during sex organ development in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **96**, 11664–11669.
- Schmid, M. W., Schmidt, A., Klostermeier, U. C., Barann, M., Rosenstiel, P. and Grossniklaus, U. (2012). A powerful method for transcriptional profiling of specific cell types in eukaryotes: laser-assisted microdissection and RNA sequencing. *PLoS ONE* **7**, e29685.
- Schmidt, A., Wüest, S. E., Vijverberg, K., Baroux, C., Kleen, D. and Grossniklaus, U. (2011). Transcriptome analysis of the *Arabidopsis* megasporophyte mother cell uncovers the importance of RNA helicases for plant germline development. *PLoS Biol.* **9**, e1001155.
- Schmidt, A., Schmid, M. W. and Grossniklaus, U. (2012). Analysis of plant germline development by high-throughput RNA profiling: technical advances and new insights. *Plant J.* **70**, 18–29.
- Schmidt, A., Schmid, M. W., Klostermeier, U. C., Qi, W., Guthörl, D., Sailer, C., Waller, M., Rosenstiel, P. and Grossniklaus, U. (2014). Apomictic and sexual germline development differ with respect to cell cycle, transcriptional, hormonal and epigenetic regulation. *PLoS Genet.* **10**, e1004476.
- Seifert, G. J. and Roberts, K. (2007). The biology of arabinogalactan proteins. *Annu. Rev. Plant Biol.* **58**, 137–161.
- Sharbel, T. F., Voigt, M. L., Corral, J. M., Thiel, T., Varshney, A., Kümlehn, J., Vogel, H. and Rotter, B. (2009). Molecular signatures of apomictic and sexual ovules in the *Boechera holboellii* complex. *Plant J.* **104**, 14026–14031.
- Sharbel, T. F., Voigt, M. L., Corral, J. M., Galla, G., Kümlehn, J., Klukas, C., Schreiber, F., Vogel, H. and Rotter, B. (2010). Apomictic and sexual ovules of *Boechera* display heterochronic global gene expression patterns. *Plant Cell* **22**, 655–671.
- She, W., Grimanelli, D., Rutowicz, K., Whitehead, M. W. J., Puzio, M., Kotlinski, M., Jerzmanowski, A. and Baroux, C. (2013). Chromatin reprogramming during the somatic-to-reproductive cell fate transition in plants. *Development* **140**, 4008–4019.
- Sheridan, W. F., Avalkina, N. A., Shamrov, I. I., Batygina, T. B. and Golubovskaya, I. N. (1996). The *mac1* gene: controlling the commitment to the meiotic pathway in maize. *Genetics* **142**, 1009–1020.
- Sheridan, W. F., Golubeva, E. A., Ahrhams, L. I. and Golubovskaya, I. N. (1999). The *mac1* mutation alters the developmental fate of the hypodermal cells and their cellular progeny in the maize anther. *Genetics* **153**, 933–941.
- Sieber, P., Petrascheck, M., Barberis, A. and Schneitz, K. (2004). Organ polarity in *Arabidopsis*. *NOZZLE* physically interacts with members of the YABBY family. *Plant Physiol.* **135**, 2172–2185.
- Singh, M., Goel, S., Meeley, R. B., Dantec, C., Parrinello, H., Michaud, C., Leblanc, O. and Grimanelli, D. (2011). Production of viable gametes without meiosis in maize deficient for an ARGONAUTE protein. *Plant Cell* **23**, 443–458.
- Spillane, C., Steimer, A. and Grossniklaus, U. (2001). Apomixis in agriculture: the quest for clonal seeds. *Sex. Plant Reprod.* **14**, 179–187.
- Spillane, C., Curtis, M. D. and Grossniklaus, U. (2004). Apomixis technology development—virgin births in farmers' fields? *Nat. Biotechnol.* **22**, 687–691.
- Sprunck, S. and Gross-Hardt, R. (2011). Nuclear behavior, cell polarity, and cell specification in the female gametophyte. *Sex. Plant Reprod.* **24**, 123–136.
- Sundaresan, V. and Alandete-Saez, M. (2010). Pattern formation in miniature: the female gametophyte of flowering plants. *Development* **137**, 179–189.
- Tucker, M. R., Araujo, A.-C. G., Paech, N. A., Hecht, V., Schmidt, E. D. L., Rossell, J.-B., de Vries, S. C. and Koltunow, A. M. G. (2003). Sexual and apomictic reproduction in *Hieracium* subgenus *pilosella* are closely interrelated developmental pathways. *Plant Cell* **15**, 1524–1537.
- Tucker, M. R., Okada, T., Hu, Y., Scholefield, A., Taylor, J. M. and Koltunow, A. M. G. (2012). Somatic small RNA pathways promote the mitotic events of megagametogenesis during female reproductive development in *Arabidopsis*. *Development* **139**, 1399–1404.
- Twill, D. (2011). Male gametogenesis and germline specification in flowering plants. *Sex. Plant Reprod.* **24**, 149–160.
- Urao, T., Miyata, S., Yamaguchi-Shinozaki, K. and Shinozaki, K. (2000). Possible His to Asp phosphorelay signaling in an *Arabidopsis* two-component system. *FEBS Lett.* **478**, 227–232.
- Vielle-Calzada, J.-P., Crane, C. F. and Stelly, D. M. (1996). Apomixis—the asexual revolution. *Science* **274**, 1322–1323.
- Völz, R., von Lyncker, L., Baumann, N., Dresselhaus, T., Sprunck, S. and Gross-Hardt, R. (2012). LACHESIS-dependent egg-cell signaling regulates the development of female gametophytic cells. *Development* **139**, 498–502.
- Wang, C.-J. R., Nan, G.-L., Kelliher, T., Timofejeva, L., Vernoud, V., Golubovskaya, I. N., Harper, L., Egger, R., Walbot, V. and Cande, W. Z. (2012). Maize *multiple archesporial cells1* (*mac1*), an ortholog of rice *TDL1A*, modulates cell proliferation and identity in early anther development. *Development* **139**, 2594–2603.
- Willemse, M. T. M. and van Went, J. L. (1984). The female gametophyte. In *Embryology of Angiosperms* (ed. B. M. Johri), pp. 159–196. Berlin: Springer-Verlag.
- Wüest, S. E., Vijverberg, K., Schmidt, A., Weiss, M., Gheyselinck, J., Lohr, M., Wellmer, F., Rahnenführer, J., von Mering, C. and Grossniklaus, U. (2010). *Arabidopsis* female gametophyte gene expression map reveals similarities between plant and animal gametes. *Curr. Biol.* **20**, 506–512.
- Wüest, S. E., Schmid, M. W. and Grossniklaus, U. (2013). Cell-specific expression profiling of rare cell types as exemplified by its impact on our understanding of female gametophyte development. *Curr. Opin. Plant Biol.* **16**, 41–49.
- Yajima, M. and Wessel, G. M. (2011). The multiple hats of Vasa: its functions in the germline and in cell cycle progression. *Mol. Reprod. Dev.* **78**, 861–867.
- Yamada, H., Suzuki, T., Terada, K., Takei, K., Ishikawa, K., Miwa, K., Yamashino, T. and Mizuno, T. (2001). The *Arabidopsis* AHK4 histidine kinase is a cytokinin-binding receptor that transduces cytokinin signals across the membrane. *Plant Cell Physiol.* **42**, 1017–1023.
- Yang, W.-C. and Sundaresan, V. (2000). Genetics of gametophyte biogenesis in *Arabidopsis*. *Curr. Opin. Plant Biol.* **3**, 53–57.

- Yang, W.-C., Ye, D., Xu, J. and Sundaresan, V. (1999). The *SPOROCTELESS* gene of *Arabidopsis* is required for initiation of sporogenesis and encodes a novel nuclear protein. *Genes Dev.* **13**, 2108-2117.
- Yang, S.-L., Xie, L.-F., Mao, H.-Z., Puah, C. S., Yang, W.-C., Jiang, L., Sundaresan, V. and Ye, D. (2003). *TAPETUM DETERMINANT1* is required for cell specialization in the *Arabidopsis* anther. *Plant Cell* **15**, 2792-2804.
- Yang, S.-L., Jiang, L., Puah, C. S., Xie, L.-F., Zhang, X.-Q., Chen, L.-Q., Yang, W.-C. and Ye, D. (2005). Overexpression of *TAPETUM DETERMINANT1* alters the cell fates in the *Arabidopsis* carpel and tapetum via genetic interaction with *excess microsporocytes1/extra sporogenous cells*. *Plant. Physiol.* **139**, 186-191.
- Youl, J. J., Bacic, A. and Oxley, D. (1998). Arabinogalactan-proteins from *Nicotiana glauca* and *Pyrus communis* contain glycosylphosphatidylinositol membrane anchors. *Proc. Natl. Acad. Sci. USA* **95**, 7921-7926.
- Zhang, D. and Yang, L. (2014). Specification of tapetum and microsporocyte cells within the anther. *Curr. Opin. Plant Biol.* **17**, 49-55.
- Zhang, Y., Yang, J. and Showalter, A. M. (2011). AtAGP18, a lysine-rich arabinogalactan protein in *Arabidopsis thaliana*, functions in plant growth and development as a putative co-receptor for signal transduction. *Plant Signal. Behav.* **6**, 855-857.
- Zhao, D.-Z., Wang, G.-F., Speal, B. and Ma, H. (2002). The *excess microsporocytes1* gene encodes a putative leucine-rich repeat receptor protein kinase that controls somatic and reproductive cell fates in the *Arabidopsis* anther. *Genes Dev.* **16**, 2021-2031.
- Zhao, X., de Palma, J., Oane, R., Gamuyao, R., Luo, M., Chaudhury, A., Hervé, P., Xue, Q. and Bennett, J. (2008). OsTDL1A binds to the LRR domain of rice receptor kinase MSP1, and is required to limit sporocyte numbers. *Plant J.* **54**, 375-387.